arXiv:1404.5559v4 [math.GT] 16 Feb 2015

Right-angled Artin groups in the C^∞ diffeomorphism group of the real line

HYUNGRYUL BAIK, SANG-HYUN KIM, AND THOMAS KOBERDA

Abstract. We prove that every right-angled Artin group embeds into the C^∞ diffeomorphism group of the real line. As a corollary, we show every limit group, and more generally every countable residually RAAG group, embeds into the C^∞ diffeomorphism group of the real line.

1. Introduction

The *right-angled Artin group* on a finite simplicial graph Γ is the following group presentation:

$$A(\Gamma) = \langle V(\Gamma) \mid [v_i, v_j] = 1 \text{ if and only if } \{v_i, v_j\} \in E(\Gamma)\rangle.$$

Here, $V(\Gamma)$ and $E(\Gamma)$ denote the vertex set and the edge set of Γ, respectively.

For a smooth oriented manifold X, we let $\mathrm{Diff}^\infty_+(X)$ denote the group of orientation preserving C^∞ diffeomorphisms on X. A group G is said to *embed* into another group H if there is an injective group homormophism $G \to H$. Our main result is the following.

Theorem 1. *Every right-angled Artin group embeds into* $\mathrm{Diff}^\infty_+(\mathbb{R})$.

Recall that a finitely generated group G is a *limit group* (or a *fully residually free group*) if for each finite set $F \subset G$, there exists a homomorphism ϕ_F from G to a nonabelian free group such that ϕ_F is an injection when restricted to F. The class of limit groups fits into a larger class of groups, which we call *residually RAAG groups*. A group G is in this class if for each $1 \neq g \in G$, there exists a graph Γ_g and a homomorphism $\phi_g \colon G \to A(\Gamma_g)$ such that $\phi_g(g) \neq 1$. Since the class of right-angled Artin groups is closed under taking finite direct products, we could replace g with an arbitrary finite subset of G. Theorem 1 can be strengthened as follows.

Corollary 2. *Every countable residually RAAG group embeds into* $\mathrm{Diff}^\infty_+(\mathbb{R})$.

A similar argument to the proof of Theorem 1 also applies to the group $\mathrm{PL}_+(\mathbb{R})$ of orientation preserving piecewise-linear homeomorphisms of $\mathbb{R}$.

Corollary 3. *Every countable residually RAAG group embeds into* $\mathrm{PL}_+(\mathbb{R})$.

Date: February 17, 2015.

Key words and phrases. right-angled Artin group, limit group, diffeomorphism group, left–greedy form.

Our construction requires infinitely many domains of linearity in $\mathbb{R}$, so that we must take infinite subdivisions of $\mathbb{R}$ in order to get residually RAAG groups inside of $\mathrm{PL}_+(\mathbb{R})$. The reader may compare Corollary 3 with the work of Brin and Squier [6], which shows that the group $\mathrm{PLF}(\mathbb{R})$ of piecewise-linear homeomorphisms of $\mathbb{R}$ with finite subdivisions contains no nonabelian free subgroup.

Recall that a group G is called *virtually special* if a finite index subgroup $H \leq G$ acts properly and cocompactly on a $\mathrm{CAT}(0)$ cube complex X such that the quotient $H\backslash X$ avoids certain pathologies of its half–planes (see [16]). A consequence of such an action is that the subgroup H embeds in a right-angled Artin group.

Corollary 4. *Let G be a group which is virtually special. Then there is a finite index subgroup $H \leq G$ which embeds into $\mathrm{Diff}^\infty_+(\mathbb{R})$ and also into $\mathrm{PL}_+(\mathbb{R})$.*

Examples of virtually special groups include fundamental groups of closed surfaces and finite volume hyperbolic 3–manifold groups [1, 22]. Combining the work of Bergeron–Haglund–Wise [3] and Bergeron–Wise [4], we have that there are virtually special closed hyperbolic manifolds in all dimensions.

The finitely presented subgroups of diffeomorphism groups are generally very complicated. In [5], Bridson used a virtually special version of the Rips machine to produce finitely presented subgroups of right-angled Artin groups with exotic algorithmic properties. An arbitrary class of groups which contains sufficiently complicated right-angled Artin groups thereby also has finitely presented subgroups with exotic algorithmic properties (cf. [18]):

Corollary 5. *Suppose $G = \mathrm{Diff}^\infty_+(\mathbb{R})$ or $G = \mathrm{PL}_+(\mathbb{R})$. Then there is a finitely presented subgroup $H \leq G$ such that the conjugacy problem in H is unsolvable. Furthermore, the isomorphism problem for the class of finitely presented subgroups of G is unsolvable.*

The reader may contrast Corollary 5 with Thompson's groups F, T, and V, in which the conjugacy problem is generally solvable [2]. The group T can be embedded into $\mathrm{Diff}^\infty_+(\mathbb{S}^1)$ [15].

The group H in Corollary 5 is not conjugacy separable, since conjugacy separable groups have solvable conjugacy problems. In [13], Farb and Franks showed that the group of real analytic diffeomorphisms of $\mathbb{R}$ contains non–solvable Baumslag–Solitar groups, which are not even residually finite.

1.1. **Notes and references.** It is well-known that right-angled Artin groups embed in $\mathrm{Homeo}(\mathbb{R})$, as follows from the fact that right-angled Artin groups admit left–invariant orders. Similarly, right-angled Artin groups embed in $\mathrm{Homeo}(\mathbb{S}^1)$ because they admit left–invariant cyclic orderings. Alternatively, one can embed an arbitrary right-angled Artin group into the mapping class group $\mathrm{Mod}(S)$ of a compact surface with one boundary, and then embed $\mathrm{Mod}(S)$ into $\mathrm{Homeo}(\mathbb{S}^1)$. As noted in [12, p.47], it is generally difficult to

smoothen these embeddings. The reader may consult [20] for a general discussion of the relationship between linear and cyclic orderings of groups and embeddings into Homeo($\mathbb{R}$) and Homeo($\mathbb{S}^1$).

Farb and Franks [14] proved that every residually torsion–free nilpotent group embeds in the group of C^1 diffeomorphisms of the interval and of the circle. This implies that residually RAAG groups embed in the group of C^1 diffeomorphisms of both the interval and of the circle. Their construction does not allow for twice differentiable diffeomorphisms. In fact, Plante and Thurston [21] showed that nilpotent groups of C^2 diffeomorphisms of the interval or of the circle are abelian. In the same vein, Farb and Franks [14] show that every nilpotent subgroup of C^2 diffeomorphisms of $\mathbb{R}$ is metabelian.

For the case when the dimension is two, Calegari and Rolfsen proved that every right-angled Artin group embeds into the piecewise linear homeomorphism group of a square fixing the boundary [7]. M. Kapovich showed that every right-angled Artin group embeds into the symplectomorphism group of the sphere [17]. The second and the third named authors refined this result by embedding every right-angled Artin group into the symplectomorphism groups of the disk and of the sphere by quasi-isometric group embeddings [18].

2. Acknowledgements

The authors thank B. Farb, A. Navas and D. Rolfsen for helpful comments. The authors are especially grateful to J. Bowden for critical comments regarding a previous version of the paper. The second named author was supported by National Research Foundation of Korean Government (NRF-2013R1A1A1058646) and Samsung Science and Technology Foundation (SSTF-BA1301-06). The third named author is partially supported by NSF grant DMS-1203964.

3. Building an injective homomorphism

Throughout this section, we let Γ be a finite graph. Consider an element g of $A(\Gamma)$. A *reduced word representing* g means a minimal length word in the standard generating set $V(\Gamma) \cup V(\Gamma)^{-1}$ representing g. The *support of* g is the set of vertices v of Γ such that v or v^{-1} appears in a reduced word representing g. We denote the support of g by $\mathrm{supp}(g)$. We say g is a *clique word* if every pair of vertices in $\mathrm{supp}(g)$ are adjacent in Γ. A *clique word decomposition for* g is the concatenation $w_k \cdots w_1$ of clique words $w_1, w_2, \ldots, w_k$ such that the concatenation is still reduced and represents g in $A(\Gamma)$. Since vertices themselves are clique words, every element in $A(\Gamma)$ has a clique word decomposition. A clique word decomposition $w_k \cdots w_1$ is *left–greedy* if for each $i < k$ and for each $v \in \mathrm{supp}(w_i)$, there exists a vertex $v' \in \mathrm{supp}(w_{i+1})$ such that $[v, v'] \neq 1$. The left–greedy clique word decomposition can be compared to the left–greedy normal form used in [19].

Lemma 6. *Every element of $A(\Gamma)$ admits a left–greedy clique word decomposition.*

Proof. Fix an element $g \in A(\Gamma)$. Let us define the *complexity* of a clique word decomposition $w_k w_{k-1} \cdots w_1$ for g as the k-tuple consisting of the word lengths $(|w_k|, |w_{k-1}|, \dots, |w_1|)$. In the lexicographical order, this complexity is bounded above by $(|g|)$. Let us assume $w_k w_{k-1} \cdots w_1$ is the maximal clique word decomposition for g in this order. If $w_k \cdots w_1$ is not left–greedy, then there exists i and $v \in \mathrm{supp}(w_{i-1})$ be such that $v \cup \mathrm{supp}(w_i)$ spans a complete subgraph of Γ. Then we can move an occurrence of v or v^{-1} in w_{i-1} to w_i, i.e. we *slide to the left*. This is a contradiction to the maximality. □

Let X be a smooth oriented manifold. If $f \in \mathrm{Diff}^\infty_+(X)$, then we write the fixed point set of f as $\mathrm{Fix}(f)$. The closure of $X \setminus \mathrm{Fix}(f)$ will be denoted as $\mathrm{supp}(f)$. For $f_1, f_2, \dots, f_n \in \mathrm{Diff}^\infty_+(X)$, let us define the *disjointness graph* Λ by $V(\Lambda) = \{v_1, v_2, \dots, v_n\}$ and

$$E(\Lambda) = \{\{v_i, v_j\} \colon i \neq j \text{ and } \mathrm{supp}(f_i) \cap \mathrm{supp}(f_j) = \varnothing\}.$$

Note that two self-maps with disjoint supports commute. Hence, we have a group homomorphism $\phi \colon A(\Lambda) \to \mathrm{Diff}^\infty_+(X)$ satisfying $\phi(v_i) = f_i$. Often, it is not an obvious task at all to decide whether or not such a map ϕ is injective; see [11, 17, 18, 7] for related work on diffeomorphism groups and [9, 10, 19, 8] on mapping class groups.

Let H be a group. Another group G is *residually H* if for each nontrivial element g in G there exists a group homomorphism $\phi_g \colon G \to H$ such that $\phi_g(g) \neq 1$. For a smooth oriented manifold X, we let $\mathrm{Diff}^\infty_+(X, \partial X)$ denote the group of orientation preserving C^∞ diffeomorphisms on X which restrict to the identity near ∂X. The crucial fact we need is the following:

Lemma 7. *The group $A(\Gamma)$ is residually $\mathrm{Diff}^\infty_+(I, \partial I)$.*

Proof. Let us fix an element g in $A(\Gamma)$. We let $w_k \cdots w_1$ be a left–greedy clique decomposition representing g. We will construct a group homomorphism

$$\phi_g \colon A(\Gamma) \to \mathrm{Diff}^\infty_+(I, \partial I)$$

such that $\phi_g(g) \neq 1$. We can inductively choose a (possibly redundant) sequence

$$v_1 \in \mathrm{supp}(w_1), v_2 \in \mathrm{supp}(w_2), \dots, v_k \in \mathrm{supp}(w_k)$$

such that $[v_i, v_{i+1}] \neq 1$ for each $i = 1, 2, \dots, k-1$. There exists $\sigma_i \in \{-1, 1\}$ and $n_i > 0$ such that $v_i^{\sigma_i n_i}$ is the highest power of v_i in w_i. This means that $v_i \notin \mathrm{supp}(w_i v_i^{-\sigma_i n_i})$.

We choose $\rho \in \mathrm{Diff}^\infty_+(\mathbb{R})$ such that $\rho(1/4) = 5/4$ and $\rho(x) = x$ for $x \leq 0$ or $x \geq 3/2$. Put $\rho_i(x) = \rho(x-i)+i$ and $I_i = [i, i+3/2]$, so that $\mathrm{supp}\, \rho_i \subseteq I_i$. Note that $I_i \cap I_j = \emptyset$ for $|i-j| > 1$.

For each $v \in V(\Gamma)$, we define

$$\psi_g(v) = \prod_{v_j = v} \rho_j^{\sigma_j}.$$

This means that if $\{j\colon v_j = v\} = \{j_1 < j_2 < \cdots < j_n\}$ then $\psi_g(v) = \rho_{j_1}^{\sigma_1}\rho_{j_2}^{\sigma_2}\cdots\rho_{j_n}^{\sigma_n}$. We use the convention that the empty multiplication is trivial. Let us write $J_v = \cup_{v_j=v} I_j \supseteq \operatorname{supp}\psi_g(v)$. Note that if $v_i = v = v_j$ and $i \neq j$, then $I_i \cap I_j = \emptyset$. For each $\{u, v\} \in E(\Gamma)$, the choice of $v_1, v_2, \ldots, v_k$ implies that $J_u \cap J_v = \emptyset$. In other words, Γ is a subgraph of the disjointness graph of $\{\psi_g(v)\colon v \in V(\Gamma)\}$. It follows that ψ_g defines a group homomorphism from $A(\Gamma)$ to $\mathrm{Diff}^\infty_+(\mathbb{R})$.

Suppose $\ell \in \{1, 2, \ldots, k\}$. For every $u \in \operatorname{supp} w_\ell \setminus \{v_\ell\}$, we have $J_u \cap J_{v_\ell} = \emptyset$. Since $\ell + 1/4 \in I_\ell \subseteq J_{v_\ell}$, we see that if $x \in [\ell + 1/4, \ell + 1/2]$ is an arbitrary point then

$$\psi_g(w_\ell)(x) = \psi_g(v_\ell^{\sigma_\ell n_\ell})(x) = \rho_\ell^{n_\ell}(x) \in [\ell + 5/4, \ell + 3/2].$$

It follows that if $y \in [5/4, 3/2]$ is arbitrary then

$$\psi_g(g)(y) = \psi_g(w_k w_{k-1} \cdots w_1)(y) \in [k + 5/4, k + 3/2].$$

In particular, $\psi_g(g)$ is not the identity. By restricting the image of ψ_g onto the interval $[0, k+2]$ and conjugating by a diffeomorphism $[0, k+2] \approx I$, we obtain a desired group homomorphism ϕ_g. □

We have the following general fact.

Lemma 8. *Let G and H be groups. If G is countable and residually H, then G embeds into the countable direct product $\prod_{\mathbb{Z}} H$.*

Proof. Define

$$\psi\colon G \to \prod_{g \in G} H$$

by $\psi(x)(g) = \phi_g(x)$, where ϕ_g is as in the definition of residually H group. □

Since $\mathrm{Diff}^\infty_+([0,\infty), \{0\}) \hookrightarrow \mathrm{Diff}^\infty_+(\mathbb{R})$, Theorem 1 a trivial consequence of the following:

Theorem 9. *Every right-angled Artin group embeds into* $\mathrm{Diff}^\infty_+([0,\infty), \{0\})$.

Proof. Immediate from Lemmas 7 and 8, as well as the fact that

$$\prod_{\mathbb{Z}} \mathrm{Diff}^\infty_+(I, \partial I) \hookrightarrow \mathrm{Diff}^\infty_+([0,\infty), \{0\}).$$ □

Proof of Corollary 2. Lemma 7 implies that a residually RAAG is residually $\mathrm{Diff}^\infty_+(I, \partial I)$. We proceed as Theorem 9. □

Proof of Corollary 3. It suffices to show that every right-angled Artin group is residually $\mathrm{PL}_+(I)$. For this, we follow the proof of Lemma 7 by using $\rho_0 \in \mathrm{PL}_+(\mathbb{R})$ defined by

$$\rho_0(x) = \begin{cases} x & \text{if } x < 0 \text{ or } x \geq \frac{3}{2} \\ 5x & \text{if } 0 \leq x < \frac{1}{4} \\ (x+6)/5 & \text{if } \frac{1}{4} \leq x < \frac{3}{2} \end{cases}$$

instead of ρ. □

References

1. Ian Agol, *The virtual Haken conjecture*, Doc. Math. **18** (2013), 1045–1087, With an appendix by Agol, Daniel Groves, and Jason Manning. MR 3104553
2. James Belk and Francesco Matucci, *Conjugacy and dynamics in Thompson's groups*, Geom. Dedicata **169** (2014), 239–261. MR 3175247
3. Nicolas Bergeron, Frédéric Haglund, and Daniel T. Wise, *Hyperplane sections in arithmetic hyperbolic manifolds*, J. Lond. Math. Soc. (2) **83** (2011), no. 2, 431–448. MR 2776645 (2012f:57037)
4. Nicolas Bergeron and Daniel T. Wise, *A boundary criterion for cubulation*, Amer. J. Math. **134** (2012), no. 3, 843–859. MR 2931226
5. Martin R. Bridson, *On the subgroups of right angled artin groups and mapping class groups*, To appear in *Math. Res. Lett.*
6. Matthew G Brin and Craig C Squier, *Groups of piecewise linear homeomorphisms of the real line*, Inventiones Mathematicae **79** (1985), no. 3, 485–498.
7. Danny Calegari and Dale Rolfsen, *Groups of PL homeomorphisms of cubes*, preprint (2014).
8. Matt T. Clay, Christopher J. Leininger, and Johanna Mangahas, *The geometry of right-angled Artin subgroups of mapping class groups*, Groups Geom. Dyn. **6** (2012), no. 2, 249–278. MR 2914860
9. John Crisp and Luis Paris, *The solution to a conjecture of Tits on the subgroup generated by the squares of the generators of an Artin group*, Invent. Math. **145** (2001), no. 1, 19–36. MR 1839284 (2002j:20069)
10. John Crisp and Bert Wiest, *Embeddings of graph braid and surface groups in right-angled Artin groups and braid groups*, Algebr. Geom. Topol. **4** (2004), 439–472. MR 2077673 (2005e:20052)
11. ———, *Quasi-isometrically embedded subgroups of braid and diffeomorphism groups*, Trans. Amer. Math. Soc. **359** (2007), no. 11, 5485–5503. MR 2327038 (2008i:20048)
12. Benson Farb, *Some problems on mapping class groups and moduli space*, Problems on mapping class groups and related topics, Proc. Sympos. Pure Math., vol. 74, Amer. Math. Soc., Providence, RI, 2006, pp. 11–55. MR 2264130 (2007h:57018)
13. Benson Farb and John Franks, *Groups of homeomorphisms of one-manifolds, I: Actions of nonlinear groups*, Preprint.
14. ———, *Groups of homeomorphisms of one-manifolds, III: Nilpotent subgroups*, Ergodic Theory Dynam. Systems **23** (2003), no. 5, 1467–1484.
15. Étienne Ghys and Vlad Sergiescu, *Sur un groupe remarquable de difféomorphismes du cercle*, Comment. Math. Helv. **62** (1987), no. 2, 185–239. MR 896095 (90c:57035)
16. Frédéric Haglund and Daniel T. Wise, *Special cube complexes*, Geom. Funct. Anal. **17** (2008), no. 5, 1551–1620. MR 2377497 (2009a:20061)
17. Michael Kapovich, *RAAGs in Ham*, Geom. Funct. Anal. **22** (2012), no. 3, 733–755. MR 2972607
18. Sang-hyun Kim and Thomas Koberda, *Anti–trees and right-angled Artin subgroups of planar braid groups*, preprint.
19. Thomas Koberda, *Right-angled Artin groups and a generalized isomorphism problem for finitely generated subgroups of mapping class groups*, Geom. Funct. Anal. **22** (2012), no. 6, 1541–1590. MR 3000498
20. Andrés Navas, *Groups of circle diffeomorphisms*, spanish ed., Chicago Lectures in Mathematics, University of Chicago Press, Chicago, IL, 2011. MR 2809110
21. Joseph Plante and William P. Thurston, *Polynomial growth in holonomy groups of foliations*, Comment. Math. Helvetici **51** (1976), no. 39, 567–584.

22. Daniel T. Wise, *From riches to raags: 3-manifolds, right-angled Artin groups, and cubical geometry*, CBMS Regional Conference Series in Mathematics, vol. 117, Published for the Conference Board of the Mathematical Sciences, Washington, DC, 2012. MR 2986461

Department of Mathematics, Cornell University, 111 Malott Hall, Ithaca, NY 14853, USA
E-mail address: hb278@cornell.edu

Department of Mathematical Sciences, Seoul National Univeristy, Seoul 151-747, Republic of Korea
E-mail address: s.kim@snu.ac.kr

Department of Mathematics, Yale University, 20 Hillhouse Ave, New Haven, CT 06520, USA
E-mail address: thomas.koberda@gmail.com